

## IMPACT OF CLIMATE CHANGE ON SUSTAINABLE MANAGEMENT OF GRAY WHALE (*ESCHRICHTIUS ROBUSTUS*) POPULATIONS: WHALE-WATCHING AND CONSERVATION

CHRISTIAN J. SALVADEO<sup>1</sup>, SALVADOR E. LLUCH-COTA<sup>2</sup>, MARTÍN O. MARAVILLA-CHÁVEZ<sup>3</sup>,  
SERGIO T. ÁLVAREZ-CASTAÑEDA<sup>2</sup>, MILENA MERCURI<sup>1</sup> and ALFREDO ORTEGA-RUBIO<sup>2\*</sup>

<sup>1</sup> Centro Interdisciplinario de Ciencias Marinas del IPN (CICIMAR), Apdo. Postal 592, La Paz, BCS 23000, Mexico

<sup>2</sup> Centro de Investigaciones Biológicas del Noroeste (CIBNOR), Apdo. Postal 128, La Paz, BCS 23000, Mexico

<sup>3</sup> Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT), Delegación Baja California Sur, Melchor Ocampo 1045, La Paz, BCS 23000, Mexico

**Abstract** - Some pods of gray whales (*Eschrichtius robustus*) breed every winter at three lagoons along the Baja California Peninsula and then migrate to feeding grounds on the Bering and Chukchi Seas. The number of whales arriving to the lagoons fluctuates yearly and is related to climate variability. We analyzed the documented climate effects on the whales over their distribution range and discuss the potential effects of global climate warming in their breeding areas. Our analysis indicates that global warming will be favorable for gray whale populations, but unfavorable for the Mexican whale-watching industry: favorable, because fluctuations in calf production have been positively correlated with the length of time that the primary feeding habitat was free of seasonal ice during the previous year. However, if gray whales change their breeding areas to northern bays, they will be exposed to new challenges, which will have repercussions on the whale-watching industry. We discuss these new challenges.


**Key words:** Baja California, breeding areas, *Eschrichtius robustus*, global warming, seasonal migration, sustainable management.

## INTRODUCTION

Pods of gray whales (*Eschrichtius robustus*, Lilljeborg, 1861) breed every winter in three lagoons along the Baja California Peninsula (Fig. 1): Laguna Ojo de Liebre, the northernmost locality, containing the largest number of whales; Laguna San Ignacio, the most preserved sanctuary for the species; and the Bahía Magdalena complex, comprising a long series of narrow lagoons behind barrier islands, which in total extends for 180 km. In these lagoons, the Mexican Government allows whale-watching activities by issuing permits to local organizations. Gray whales have entered some of the lagoons inside the Gulf of

California, Bahía Santa María in the State of Sinaloa and Yavaros in the State of Sonora (Figure 1), which have been abandoned (Findley and Vidal, 2002).

At the end of the breeding season, the whales migrate to the feeding grounds in the Bering and Chukchi Seas (Fig. 1), where they feed on benthic fauna. The population of gray whales seems to have reached carrying capacity, with the population size fluctuating between 20,000 and 22,000 (Rugh et al., 2008). Only one third of this population, mainly pregnant females, mature females, males, and some juveniles, travel south in late fall to participate in calving and mating. The rest of the population spreads out along



**Figure 1.** Localities of gray whale breeding and feeding along the eastern Pacific; Gulf of California (GC).

the coast of North America, from Alaska to California (Urbán et al., 2003a).

Despite yearly variations, the number of whales arriving at the lagoons seems to follow a pattern that is apparently connected to climate variations. We have reviewed the documented climatic effects on these whales over their range and herein address the potential effects of global climate warming (GCW) on their use of traditional breeding areas, and examine the impact on the whale-watching industry.

#### *Biology, migratory patterns, and habitats*

The gray whale is the only member of the Eschrichtiidae. These are medium-sized baleen whales, usually grayish-black when born; they develop scars or white spots as they grow. They do not show a true dorsal fin; instead, a series of knobs or small humps, usually 10 to 15, appear on the final third of the back. They are host to more external parasites and epizooties than any other cetacean. This includes barnacles that eventually form large colonies embedded in the skin and three species of whale lice (Cyamids) that feed on skin and damaged tissue (Jones and Swarts,

2002). Gray whales reach 14 m in length and weigh 45 metric tons. Adult females on average are larger than males. When born, calves are about 4.5 m long. Females nurse their young for 8 to 9 months. Females become sexually mature at 8 years (range is 5-11 years) and usually give birth to a single calf every other year. There seems to be no long-term bonds between males and females or any bonds between males and newborns (Rice and Wolman, 1971).

Gray whales feed primarily on small benthic organisms and this may restrict their distribution to coastal environments in the Chukchi and Bering Seas (Fig. 1), where they feed on amphipods as well as mysids, mollusks, polychaetes, and hydrozoa (Rice and Wolman, 1971). When traveling, gray whales tend to be gregarious, but are not highly social. Larger aggregations in tens or even thousands can occur in a particularly rich feeding area. In the breeding areas, large aggregations of mothers with young and courting/mating whales are common (Urbán et al., 2003a).

Only one-third of the estimated northeastern population makes the longest well-known migration

southward (a 16,000 km round trip). Mature females and males, and some juveniles of both sexes, reach the warm quiet waters in the coastal lagoons along the east side of the Baja California Peninsula (Fig. 1) (Poole, 1984). Each calf stays with its mother for feeding. Mother and calf remain active inside the lagoons and start migration when the calf is about two months old. The gray whale is the only species to bear young in warm, shallow, coastal areas and lagoons (Berta et al, 2006; Jones and Swartz 2002).

Courting animals usually stay close to the entrances of the lagoons. Mature males and females may collect in small groups of usually of one female and two males, forming a “courting triad”. Estrous females copulate with several males during the season. The estrous period lasts about three weeks near the end of December. Impregnation usually occurs during the first ovulation. If fertilization does not occur at that time, a second estrous period occurs 40 days later. An ideal reproductive cycle takes two years. Males reach maturity when they are about 11 m long and females reach maturity when they are about 11.7 m. Sexual maturity occurs between 5 and 11 years (Berta et al., 2006; Rice and Wolman, 1971).

## MATERIAL AND METHODS

We reviewed and analyzed all the available information, even in non-published papers, on the current protection policy, the whale-watching industry and the documented climate change impacts on marine mammals and gray whale populations.

## RESULTS

### *Current protection policy (Mexico and US) and population status*

In the 18th century, American whalers severely depleted the population, using the winter breeding concentrations in Mexican waters because their meat and blubber was a valuable resource. By 1900, the gray whale was near extinction. Gray whales were first protected by Mexican law at the beginning of 20th century, later by the League of Nations,

and now by the International Whaling Commission. A few native peoples of North America and Russia continue to hunt them in small numbers. The impact of aboriginal whaling was relatively small compared with the effects of industrial whaling (Reilly et al., 2008).

With no commercial hunting for over 50 years, the species has recovered to close to its pre-exploitation numbers of about 20,000 (Rugh et al., 2008). They seem to have reached carrying capacity, and the population now fluctuates with food supply (Moore et al., 2003). Gray whales were listed as Endangered under the U.S. Endangered Species Conservation Act in 1969. Further protection was given by the Marine Mammal Protection Act in 1972 and the U.S. Species Act of 1973. In 1994, the species was removed from the List of Endangered and Threatened Wildlife and Plants, when the population reached 21,000. It has also been down-listed in the World Conservation Union's 1996 IUCN Red List of Threatened Animals (from endangered to conservation dependent).

In 1972, the first action to protect individuals in their breeding areas was undertaken by the Mexican Government with the establishing of the first marine sanctuary for whales. In July 1972, the first lagoon system (Ojo de Liebre, Guerrero Negro, and Manuella) became a protected area for whales and their newborn (Diario Oficial, 1972). In 1979, Laguna San Ignacio received the same status (Diario Oficial, 1979). Since then, these two areas have been under strict Federal regulations for whale-watching activities. In 1988, the Government established the largest Biosphere Reserve in Latin America (Vizcaino Biosphere Reserve). This UNESCO World Heritage Site provided sanctuaries for the gray whale and other animals. Other winter visitors to Mexican waters include the blue whale (*Balaenoptera musculus*, Linnaeus, 1758) and the humpback whale (*Megaptera novaeangliae*, Borowski, 1781). In May 2002, the Mexican Government authorities created the Marine National Sanctuary for whales (Diario Oficial, 2002), which included the Economic Exclusion Zone (to 200 nautical miles). With this effort, more than

**Table 1.** Gray whale responses to climate variability.

Climate events (scales)	Whales response	References
ENSO (Interannual)	Distribution shift in breeding areas (northernmost distribution during El Niño & southernmost distribution during La Niña conditions).	(Gardner & Chávez, 2000) (Urban et al., 2003b)
	Increased mortality along their distribution, by starvation resulting from a reduction in prey availability in their feeding areas.	(Le Boeuf et al., 2000) (Moore et al., 2003) (Cardenas, 2004)
	Shift in the distribution of their feeding areas and diet.	(Moore et al., 2007; Moore, 2008)
Ice extension (interannual)	Less (more) ice in feeding areas, implies a longer (shorter) feeding season, which results in more (less) births.	(Perryman et al., 2002)
Regimen shift 76-77 (interdecadal)	Shift in timing migration.	(Rugh et al., 2001)
Global Climate Warming (secular or more)	Less whales in the Gulf of California.	(Table 2)
	Increased numbers of mothers with calf in California coast.	Shelden et al., (2004).
	Winter occurrence of whales on their feeding areas.	(Stafford et al., 2007; Moore and Huntington, 2008)
	Recolonization of the Atlantic Ocean by gray whales.	<a href="http://news.sciencemag.org/sciencenow/2010/05/scienceshot-are-gray-whales-retu.html">http://news.sciencemag.org/sciencenow/2010/05/scienceshot-are-gray-whales-retu.html</a>
	Decrease in whale numbers in the breeding lagoons.	(Urbán et al., 2010)

**Table 2.** Evidence of a major use of the northern Gulf of California in the past by this species.

Date	Record	References
1854 to 1874	11 whales per year were killed	Henderson, 1984
1979 to 1989	5 whales per year were sighted	Tershy and Breese, 1991 Silber et al., 1994 Sanchez-Pacheco et al., 2001
Actually	sightings are rare or even null	Heckel, 2006 Cárdenas Hinojosa pers. Comm.

45 species of cetaceans living or visiting these waters were protected.

*The whale-watching industry along the west coast of Baja California*

The past damage caused by whaling, the recovery of the species, the pristine condition of the environment and the surroundings of the breeding lagoons have been important factors in the development of whale-watching on the Baja California Peninsula. This activity began in 1972 in Laguna San Ignacio,

expanding to the other two lagoons due to increased demand, which increased from less than 1,000 visitors in the early 80s to more than 28,000 during the late 90s, becoming, with fisheries, the most important activity for these areas (Dedina and Young, 1995; Heckel et al., 2003; Sánchez-Pacheco, 1998).

Besides gray whales, in order to protect other species of resident or migratory cetaceans, the Mexican Government has established regulations and rules of behavior for whale-watching activities (NOM-131-SEMARNAT-1998) (Diario Oficial, 2000). Regula-

tions involve limiting the number of permits for boat owners engaged in whale watching in specific areas. In addition, boat operators are obliged to assist in and take courses in regulations and environment protection taught by experts in both regulations and whale behavior. These regulations have controlled the number of visitors and boats that can surround the whales.

For whale watching, observation zones were established for each lagoon, limiting access to the nursing and birthing areas. The number of boats moving at the same time inside the zones and the number of people onboard is strictly regulated, as are distances and ways to approach individuals or groups of whales. Boat operators, who are local residents, approach and maneuver among concentrations of whales with a caution learned from years of experience.

#### *Documented climate change impacts on marine mammals and gray whale*

Climate affects all components of marine ecosystems. Physical changes linked to potential future climate scenarios include reduced sea ice, rising sea levels, and changes in the distribution of preferred temperature water masses. These processes could affect marine mammal populations through reproductive success, changing migration patterns, phonological disruptions, and food availability due to changes in abundance and community structure of their prey at specific locations (Learmonth et al., 2006). This sort of impacts have already been documented as responses to short-term climate variations, such as El Niño (e.g. Majluf and Reyes, 1989; Shane, 1995; Tershy et al., 1991; Trillmich and Limberger, 1985; Tynan, 1999; Wells et al., 1990).

For lower frequency variations (i.e. decadal), abundance (Springer, 1999) and social behavior (Lusseau et al., 2004) changes have been related to fluctuations in the availability of food. The well-documented mid-1970s climate shift affected the distribution and abundance of several ecosystem components, such as salmon (*Oncorhynchus* spp., Walbaum,

1792), hake (*Merluccius productus*, Ayres, 1855), sardines (*Sardinops sagax*, Jenyns, 1842) and several groundfish species (McFarlane et al., 2000). Similarly, marine mammals responded to these ecosystem changes (Trites et al., 2007; Wade et al., 2007).

Northern hemisphere warming during the last decades has been linked to a large decline in marine mammal abundance, suggesting that in spite of their migratory capacity, they are also vulnerable to climate change (MacLeod et al., 2005; Moore, 2008, Salvadeo et al., 2010; Stirling et al., 1999; Tynan & DeMaster 1997). Particularly for gray whales, Table 1 summarizes different evidences of climate change impacts at different scales, affected mainly their distribution, reproduction, survival and timing migration.

## DISCUSSION

### *Potential future scenarios of gray whale abundance and distribution*

MacLeod (2009) recently provided a useful framework for assessing which cetacean species' ranges are likely to change as a result of increases in water temperature and whether they will expand, shift poleward, or contract based on their current distributions, and the variety of the effects on feeding (polar regions) and breeding areas (temperate and subtropical regions), and changes in the migratory patterns.

Particularly for gray whale, a poleward shift is likely occurring: a) during El Niño years they tend to use northern areas more intensively than in normal years, while during La Niña they tend to use southern regions (Gardner and Chávez-Rosales, 2000; Urbán et al., 2003b); b) there is an increase in calf sightings at northern stations that correlates with warmer sea surface temperature anomalies (Shelden et al., 2004); c) there seems to be a range expansion into Arctic waters (Moore and Huntington, 2008); d) within the Mexican lagoons, there is an apparent long-term (century) tendency in the use of breeding lagoons, increasing in the northern (Ojo de Liebre) and decreasing in the southern (Bahía Magdalena; Urbán et



al., 2003a); e) there was a decrease in the numbers of whales in the breeding lagoons during the seasons of 2007 to 2010, also observed from shore-based surveys at Piedras Blancas during the northbound migration (Urbán et al., 2010); f) the unusual sighting of a gray whale in the Mediterranean Sea is another possible effect of their expansion to the north, which allows them to cross the Arctic to the Atlantic (<http://news.sciencemag.org/sciencenow/2010/05/scienceshot-are-gray-whales-retu.html>); and g) in spite of having an increasing population of gray whales in the eastern Pacific, the observations of individuals inside the Gulf of California have been consistently declining (Table 2).

It should be noted that Findley and Vidal (2002) described a decline in the presence of gray whales in the continental Gulf coast and related this to increases in human activities. But, we believe that GCW is the main cause of their decline inside the Gulf of California. Our suggestion is based on: a) gray whales migrate even more to the south than Bahía Magdalena during cooler winters (La Niña conditions) and in some cases cross to the continental coast; and b) there have been only six documented incidents of gray whales entangled in passive fishing gear (none of them inside the Gulf of California), and there are no records of strikes by ships on gray whales in Mexico (Urbán et al., 2003a).

#### *Challenges for a sustainable ecotourism industry and gray whale conservation efforts*

After reviewing and analyzing the information presented, we propose that GCW will be potentially favorable for the gray whale population and certainly unfavorable for the Mexican whale-watching industry: potentially favorable for the species because it has already been documented that years with extensive seasonal ice reduce the length of time the primary feeding habitat is available for pregnant females, which consequently results in a suboptimal nutritive condition and a low calf production (Perryman et al., 2002). Therefore, less sea ice extension implies a longer time and a larger area for feeding, favoring a better body condition and calf

production. However, we consider that the effect of GCW on the gray whale population is only “potentially” favorable, because if gray whales change their breeding areas to northern bays, they will be exposed to new problems. The northern bays are more populated and more open than the coastal lagoons in Baja California Sur. There will be challenges for protecting them in the more open seas near heavily populated urban areas, and authorities and conservation managers are not currently prepared for such shift. Further studies will be necessary in the north of the Baja California Lagoons (especially during El Niño warmer years) to evaluate the potential problems of these northern areas. In particular, a description of the geographic and oceanographic characteristics will be required, with special emphasis on the identification of potential problems generated by the overlapping with human activities such as offshore oil and gas development, commercial fishing, vessel traffic, whale watching, scientific research and coastal tuna farms (e.g. Heckel et al., 2001, 2003; Moore and Clarke, 2002).

If these potential problems are well identified and characterized, managers will be allowed to work on bringing in new legislation and regulations to suit the new reality, taking into account all the new actors involved, including new whale-watching tour operators, thus contributing to the sustainable management of this species in these northern areas.

Special attention must be devoted to the southern Mexican coastal zones. Particularly for the Mexican Baja California Sur whale-watching industry, this range shift will certainly be unfavorable; there will be a significant decline in gray whale arriving numbers to these coastal lagoons. The main problem with the ecotourism activities in these zones is that it focuses exclusively on gray whale. The social effects of the decrease in this whale-watching industry will be devastating for the precarious economy of these small towns and it must be assessed. Considering the existing social infrastructure related to ecotourism around these Mexican coastal lagoons and the pristine and biological diversity of the area, it is mandatory to initiate a search for alternative ecotourism

activities that can contribute to the sustainable development of these regions.

**Acknowledgments** - We thank the Consejo Nacional de Ciencia y Tecnología (CONACyT) and the Programa Institucional de Formación de Investigadores from the Instituto Politécnico Nacional (PIFI-IPN) for the scholarships given to C.S. Financial support was provided by CONACYT-REDES project 2012-NOO23-194637. Ira Fogel of CIBNOR provided valuable editorial comments.

## REFERENCES

- Berta, A., Sumich, J. L. and K. M. Kovacs (2006). Marine mammals evolutionary biology. San Diego: Academic Press.
- Cárdenas-Hinojosa, G. (2004). Análisis histórico de la mortalidad de ballena gris (*Eschrichtius robustus*) en la costa occidental de la Península de Baja California, México. BSc thesis, Universidad Autónoma de Baja California Sur.
- Dedina, S. and E. Young (1995). Conservation and development in the gray whale lagoons of Baja California Sur, México. *U.S. Marine Mammal Comisión Report*. Washington D.C.
- Diario Oficial (1972). Decreto que declara zona de refugio para ballenas y ballenatos, las aguas del área de la Laguna Ojo de Liebre, al sur de la Bahía de Sebastián Vizcaíno, en el Litoral del Océano Pacífico, Territorio de Baja California. *Diario Oficial de la Federación* 14 de enero de 1972: 4-5.
- Diario Oficial (1979). Decreto por el que se declara como refugio para ballenas grávidas y ballenatos y zona de atracción turístico-marítima, las aguas de la zona interior de la Laguna de San Ignacio, en el Estado de Baja California Sur. *Diario Oficial de la Federación* 16 de Julio de 1979: 23-4.
- Diario Oficial (1988). Decreto por el que se declara la Reserva de la Biósfera 'El Vizcaíno', ubicada en el municipio de Mulegé, B.C.S. *Diario Oficial de la Federación* 30 de noviembre de 1988: 2-26.
- Diario Oficial (2000). Norma Oficial Mexicana NOM-131-SEMARNAT-1998. Que establece lineamientos y especificaciones para el desarrollo de actividades de observación de ballenas, relativas a su protección y a la conservación de su hábitat, Mexico City. *Diario Oficial de la Federación* 10 enero de 2000: 11-7.
- Diario Oficial (2002). Acuerdo por el que se establece como área de refugio para proteger a las especies de grandes ballenas de los subórdenes Mysticeti y Odontoceti, las zonas marinas que forman parte del territorio nacional y aquellas sobre las que la nación ejerce su soberanía y jurisdicción. *Diario Oficial de la Federación* 24 de mayo de 2002.
- Findley, L. T. and O. Vidal (2002). Gray whale (*Eschrichtius robustus*) at calving sites in the Gulf of California, México. *J. Cet. Res. and Mang.* **4**, 27-40.
- Gardner, S. C., and S. Chávez-Rosales (2000). Changes in the relative abundance and distribution of gray whales (*Eschrichtius robustus*) in Magdalena Bay, Mexico during an El Niño event. *Mar. Mamm. Sci.* **16**, 728-738.
- Heckel, G. (2006). *Diversidad, distribución y abundancia de cetáceos en Bahía de los Ángeles y Canal de Ballenas, Golfo de California: bases científicas para una nueva área de observación turística de mamíferos marinos*. SNIB-CONABIO project No.DE008, final report, Mexico City.
- Heckel, G., Reilly, S. B., Sumich, J. L. and I. Espejel (2001). The influence of whale-watching on the behaviour of migrating gray whales (*Eschrichtius robustus*) in Todos Santos Bay and surrounding waters, Baja California, Mexico. *J. Cet. Res. and Mang.* **3**, 227-237.
- Heckel, G., Espejel I. and D. Fischer (2003). Issue definition and planning for whale-watching management strategies in Ensenada, Mexico. *Coas. Manag.* **31**, 277-296.
- Henderson, D. A. (1984). Nineteenth century gray whaling: grounds, catches and kills, practices and depletions of the whale population. In: M.L. Jones, S.L. Swartz, & S. Leatherwood (Eds.), *The gray whale, Eschrichtius robustus* pp. 159-186. New York: Academic Press.
- Jones, M. L., and S. L. Swartz (2002). Gray whale *Eschrichtius robustus*. In: W. F. Perrin, B. Würsig, & J. G. M. Thewissen (Eds.), *Encyclopedia of Marine Mammals* pp. 524-536. San Diego: Academic Press.
- Le Boeuf, B. J., Pérez-Cortés, M. H., Urbán, R. J., Mate, B. R., and U. F. Ollervides. (2000). High gray whale mortality and low recruitment in 1999: potential causes and implications. *J. Cet. Res. and Mang.* **2**, 85-99.
- Learmonth, J. A., Santos, M. B., Pierce, G. J., Crick, H. Q. P., and R. A. Robinson (2006). Potential effects of climate change on marine mammals. *Ocean. Mar. Biol. Ann. Rev.* **44**, 431-464.
- Lusseau, D., Willians, R., Wilson, B., Grellier, K., Hammond, P. S., and P. M. Thompson (2004). Parallel influence of climate on the behaviour of Pacific killer whales and Atlantic bottlenose dolphins. *Ecol. Lett.* **7**, 1068-1076.
- MacLeod, C. D., Bannon, S. M., Pierce, G. J., Schweder, C. S., Learmonth, J. A., Herman, J. S., and R. J. Reid (2005). Climate change and the cetacean community of north-west Scotland. *Biol. Cons.* **124**, 477-483.
- MacLeod, C. D. (2009). Global climate change, range changes and potential implications for the conservation of marine cetaceans: a review and synthesis. *Endan. Sp. Res.* **7**, 125-136.

- Majluf, P., and J. C. Reyes (1989). The marine mammals of Peru: a review. In: D. Pauly, P. Muck, J. Mendo & I. Tsukayama (Eds.), *The Peruvian upwelling system: dynamics and interactions* pp. 344-363. ICLARM Conf Proc 18.
- McFarlane, G. A., King, J. R., and R. J. Beamish (2000). Have there been recent changes in climate? Ask the fish. *Progress in Oceanography*, **47**, 147-169.
- Moore, S. E. (2008). Marine mammals as ecosystem sentinels. *J. Mamm.* **89**, 534-540.
- Moore, S. E. and J. T. Clarke (2002). Potential impact of offshore human activities on gray whales (*Eschrichtius robustus*). *J. Cet. Res. and Mang.* **4**, 19-25.
- Moore, S. E., Grebmeier, J. M., and J. R. Davies (2003). Gray whale distribution relative to forage habitat in the northern Bering Sea: current conditions and retrospective summary. *Can. J. Zool.* **81**, 734-742.
- Moore, S. E., and H. P. Huntington (2008). Arctic marine mammals and climate change: impacts and resilience. *Ecol. Appl.* **18**, 157-165.
- Moore, S. E., Wynne, K. M., Clement-Kinney, J., and J. M. Grebmeier (2007). Gray whale occurrence and forage southeast of Kodiak Island, Alaska. *Mar. Mamm. Sci.* **23**, 419-428.
- Perryman, W. L., Donahue, M. A., Perkins, P. C., and S. B. Reilly (2002). Gray whale calf production 1994-2000: Are observed fluctuations related to changes in seasonal ice cover? *Mar. Mamm. Sci.* **18**, 121-144.
- Poole, M. M. (1984). Migrations corridors of gray whales along the central California coast 1980-1982. In: M. L. Jones, S. L. Swartz, & S. Leatherwood (Eds.), *The gray whale, Eschrichtius robustus* pp. 389-408. New York: Academic Press.
- Reilly, S. B., Bannister, J. L., Best, P. B., Brown, M., Brownell Jr., R. L., Butterworth, D. S., Clapham, P. J., Cooke, J., Donovan, G. P., Urbán, J. and A. N. Zerbini (2008). *Eschrichtius robustus*. In: IUCN 2010: IUCN Red List of Threatened Species. Version 2010.2.
- Rice, D. W., and A. A. Wolman (1971). *The life history and ecology of the gray whale (Eschrichtius robustus)*. American Society of Mammalogists, Special Publication No. 3 Stillwater, Oklahoma.
- Rugh, D. J., Breiwick, J., Muto, M. M., Hobbs, R. C., Sheldon, K. W., D'Vincent, C., Laursen, I. M., Rief, S. L., Maher, S. L., and S. D. Nilson (2008). Report of the 2006-2007 census of the eastern North Pacific stock of gray whales. Alaska Fisheries Science Center. Seattle. 157 p.
- Rugh, D. J., Sheldon, K. E. W., and A. Schulman-Janiger (2001). Timing of the gray whale southbound migration. *J. Cet. Res. and Mang.* **3**, 31-39.
- Salvadeo, C. J., Lluch-Belda, D., Gómez-Gallardo, A., Urbán-Ramírez, J. and C. D. MacLeod (2010). Climate change and a poleward shift in the distribution of the Pacific white-sided dolphin in the northeastern Pacific. *Endan. Sp. Res.* **11**, 13-19.
- Sánchez-Pacheco, J. A. (1998). The tourist observation of gray whales in Baja California Sur: description and economic value. *Gac. Ecol.* **44**, 45-50.
- Sánchez-Pacheco, J. A., Vázquez Hanckin, A., and R. De Silva Davila (2001). Gray whales' mid-spring feeding at Bahía de los Angeles, Gulf of California, Mexico. *Mar. Mamm. Sci.* **17**, 186-191.
- Silber, G. K., Newcomer, M. W., Silber, P. C., Pérez Cortés, H., and G. M. Ellis (1994). Cetaceans of the northern Gulf of California - distribution, occurrence, and relative abundance. *Mar. Mamm. Sci.* **10**, 283-298.
- Shane, S. H. (1995). Relationship between pilot whales and Risso's dolphins at Santa Catalina Island, California, USA. *Mar. Ecol. Prog. Ser.* **123**, 5-11.
- Shelden, K. W., Rugh, D. J., and A. Schulman-Janiger (2004). Gray whales born north of Mexico: indicator of recovery or consequence of regime shift? *Ecol. Appl.* **14**, 1789-1805.
- Springer, A. M. (1999). Is it all climate change? Why marine bird and mammal populations fluctuate in the North Pacific. In: G. Holloway, P. Muller & D. Henderson (Eds.), *Proceedings Aha Huliko'a, Hawaiian Winter Workshop*, Manoa, University of Hawaii.
- Stafford, K. M., Moore, S.E., Spillane, M., and S. Wiggins (2007). Gray whale calls recorded near Barrow, Alaska throughout the winter of 2003-04. *Arctic.* **60**, 167-172.
- Stirling, I., Lunn, N. J., and J. Lacoza (1999). Long-term trends in the population ecology of polar bears in western Hudson Bay in relation to climatic change. *Arctic.* **52**, 294-306.
- Stirling, I. (2002). Polar bears and seals in the eastern Beaufort Sea and Amundsen Gulf: A synthesis of population trends and ecological relationships over three decades. *Arctic.* **55**, 59-76.
- Tershy, B., and D. Breese (1991). Sightings and feeding of gray whales in the northern Gulf of California. *J. Mamm.* **72**, 830-831.
- Tershy, B. R., Breese, D., and S. Álvarez-Borrego (1991). Increase in cetacean and seabird numbers in the Canal de Ballenas during an El Niño-Southern Oscillation event. *Mar. Ecol. Prog. Ser.* **69**, 299-302.
- Trillmich, F., and D. Limberger (1985). Drastic effects of El Niño on Galapagos pinnipeds. *Ecologia.* **67**, 19-22.
- Trites, A. W., Deecke, V. B., Gregr, E. J., Ford, J. K. B., and P.F. Oleisiuk (2007). Killer whales, whaling and sequential mega-



- faunal collapse in the North Pacific: a comparative analysis of dynamics of marine mammals in Alaska and British Columbia following commercial whaling. *Mar. Mamm. Sci.* **23**, 751-765.
- Tynan, C. T. (1999). Redistributions of cetaceans in the southeast Bering Sea relative to anomalous oceanographic conditions during the 1997 El Niño. *PICES Science Report*, **10**, 115-117.
- Tynan, C. T., and D. P. DeMaster (1997). Observations and predictions of Arctic climate change: potential effects on marine mammals. *Arctic*. **50**, 308-322.
- Urbán, R. J., Rojas, L. B., Pérez-Cortéz, H., Gómez-Gallardo, A. U., Swartz, S. Ludwig, S. and L. Brownell (2003a). A review of gray whales on their wintering grounds in Mexican waters. *J. Cet. Res. and Mang.* **5**, 281-295.
- Urbán, R. J., Gómez-Gallardo, U. A. and S. Ludwig (2003b). Abundance and mortality of gray whales at Laguna San Ignacio, Mexico, during the 1997-98 El Niño and the 1998-99 La Niña. *Geof. Internl.* **42**, 439-446.
- Urbán, R. J., Gómez-Gallardo U. A., Rojas-Bracho, L. and S. L. Swartz (2010). Historical changes of gray whales abundance in San Ignacio and Ojo de Liebre breeding lagoons, Mexico. *IWC Scientific Committee Paper*, SC/62/BRG36, 11 pp.
- Wade, P. R., Burkanov, V. N., Dahlheim, M. E., Friday, N. A., Fritz, L. W., Loughlin, T. R., Mizroch, S. A., Muto, M. M., Rice, D. W., Barrett-Lennard, L. G., Black, N. A. N., Burdin, A., Calambokidis, J., Cerchio, S., Ijford, J. K. B., Jjacobsen, J. K., Matkin, C. O., Matkin, N., Mehta, A. V., Small, R. J., Straley, J. M., McCluskey, S. M., Vanblaricon, G. R., and P. J. Clapham (2007). Killer whales and marine mammal trends in the North Pacific, a re-examination of evidence for sequential megafauna collapse and the prey-switching hypothesis. *Mar. Mamm. Sci.* **23**, 766-802.
- Wells, R. S., Hansen, L. J., Baldridge, A., Dohl, T. P., Kelly, D. L., and R. H. DeFran (1990). Northward extension of the range of bottlenose dolphins along the California coast. In: Leatherwood & R.R. Reeves (Eds.), *The Bottlenose Dolphin* pp. 421-431. San Diego: Academic Press.

